


Disk-driven galaxy transformation at $z \sim 4$: insights from spatially resolved ALMA data

Takafumi Tsukui (Australian National University)



**Australian
National
University**

Quenched galaxies already exist less than 2 Gyrs from Big Bang.




Madau+14



Quenched compact massive galaxies ($10^{10} - 10^{11} M_{\odot}$) at $z=3$ to 4 identified by JWST (e.g., Ito+23, Pérez-González+22)


$z \sim 2$ quenched galaxies show rotating disk (e.g., Newman+18, Toft+17).




Newman+18

Submillimeter galaxies resolved by ALMA


Dusty starburst galaxies form stars at most $> 1000 M_{\odot} \text{ yr}^{-1}$, forming the majority of stars of a massive nearby galaxy in \sim a few 100 Myr (Narayanan+15)



Simpson et al. 2020




SFR $\sim 100 M_{\odot} \text{ yr}^{-1}$
Gas depletion time = $\sim 360 \text{ Myr}$




Rizzo+2020

SFR $\sim 352 M_{\odot} \text{ yr}^{-1}$
Gas depletion time $\sim 40 \text{ Myr}$



Lelli+2021

SFR $\sim 1000 M_{\odot} \text{ yr}^{-1}$
Gas depletion time $\sim 20 \text{ Myr}$




Tsukui and Iguchi 2021

SFR $\sim 5000 M_{\odot} \text{ yr}^{-1} (?)$
Gas depletion time $\sim 20 \text{ Myr}$

Evolutionally link between two populations


Dusty star-forming galaxies might be an ancestor of quenched galaxies $z > 2$
(e.g., Barro+2013, Toft+2014)



The simple empirical model bridges the two populations.


Long+22

Evolutionally link between two populations



SFR $\sim 100 M_{\odot}\text{yr}^{-1}$


Gas depletion time = $\sim 360\text{Myr}$



SFR $\sim 352 M_{\odot}\text{yr}^{-1}$

Gas depletion time $\sim 40\text{Myr}$


Rizzo+2020



Lelli+2021

SFR $\sim 1000 M_{\odot}\text{yr}^{-1}$

Gas depletion time $\sim 20\text{Myr}$




Tsukui and Iguchi 2021

SFR $\sim 5000 M_{\odot}\text{yr}^{-1}(?)$

Gas depletion time $\sim 20\text{Myr}$

Massive compact quiescent?

Newman+17




Ito+23

Secular evolution?

Extended quenched spiral?


Fudamoto +2022



どのような条件でスターバーストが起き、遷移の過程で何が起きているか?


ALMA: ダストを見通し、空間分解した 1) 銀河構造、ダイナミクス and 2) 正確な星形成分布の把握

Evolutionally link between two populations



SFR $\sim 100 M_{\odot}\text{yr}^{-1}$


Gas depletion time = $\sim 360\text{Myr}$




SFR $\sim 352 M_{\odot}\text{yr}^{-1}$

Gas depletion time $\sim 40\text{Myr}$

Rizzo+2020




Secular evolution?



Lelli+2021

SFR $\sim 1000 M_{\odot}\text{yr}^{-1}$

Gas depletion time $\sim 20\text{Myr}$




Tsukui and Iguchi 2021

SFR $\sim 5000 M_{\odot}\text{yr}^{-1}(?)$

Gas depletion time $\sim 20\text{Myr}$

Massive compact quiescent?


Newman+17



Ito+23

Extended quenched spiral?

Fudamoto +2022





どのような条件でスターバーストが起き、遷移の過程で何が起きているか?

ALMA: ダストを見通し、空間分解した 1) 銀河構造、ダイナミクス and 2) 正確な星形成分布の把握

Evolutionally link between two populations

Massive compact quiescent?

Newman+17




Rizzo+2020

SFR $\sim 100 M_{\odot} \text{yr}^{-1}$

Gas depletion time = $\sim 360 \text{Myr}$


SFR $\sim 352 M_{\odot} \text{yr}^{-1}$

Gas depletion time $\sim 40 \text{Myr}$



Ito+23


Secular evolution?



Lelli+2021

SFR $\sim 1000 M_{\odot} \text{yr}^{-1}$

Gas depletion time $\sim 20 \text{Myr}$




Tsukui and Iguchi 2021

SFR $\sim 5000 M_{\odot} \text{yr}^{-1} (?)$

Gas depletion time $\sim 20 \text{Myr}$

Extended quenched spiral?

Fudamoto +2022



どのような条件で爆発的星形成が起き、遷移の過程で何が起きているか?

ALMA: ダストを見通し、空間分解した 1) 銀河構造、ダイナミクス and 2) 正確な星形成分布の把握


BRI 1335-0417 (z=4.41)

- Quasar-host hyper luminous infrared galaxy $L_{\text{FIR}} = 3.1 \times 10^{13} L_{\odot}$ (Carilli et al. 1998)
- $\text{SFR} = 5000 M_{\odot} \text{yr}^{-1}$ (Dust SED AGNを考慮に入っていない: Wagg et al. 2014)

Tsukui and Iguchi 2021


Tsukui + 2023c in press

VLA CO (2-1) velocity map



(Riechers+08)



ALMA [CII] observation



- VLAでは不規則な形態と力学 高い星形成率と合わせてmajor mergerだと思われていた。
- ALMAは規則的な円盤($V/\sigma \sim 2.6$) 渦巻き構造とバー構造。

BRI 1335-0417 (z=4.41)

- Quasar-host hyper luminous infrared galaxy $L_{\text{FIR}} = 3.1 \times 10^{13} L_{\odot}$ (Carilli et al. 1998)
- $\text{SFR} = 5000 M_{\odot} \text{yr}^{-1}$ (Dust SED AGNを考慮に入れていない: Wagg et al. 2014)




Tsukui and Iguchi +21

Tsukui + 2023c in press

- [CII]輝線力学モデルはバルジの存在を示唆 ($5.2 \times 10^9 M_{\odot} < 1.3 \text{kpc}$)
- 力学質量 ~ CO輝線から求めた分子ガス質量 -> gas-rich baryon-dominated disk ($f_{\text{gas}} \sim > 73\%$).

BRI 1335-0417 ($z=4.41$)


- Quasar-host hyper luminous infrared galaxy $L_{\text{FIR}} = 3.1 \times 10^{13} L_{\odot}$ (Carilli et al. 1998)
- $\text{SFR} = 5000 M_{\odot} \text{yr}^{-1}$ (Dust SED AGNを考慮に入れていない: Wagg et al. 2014)



Tsukui and Iguchi +21

- Toomre Q parameterの観点から円盤全体でガス円盤は重力不安定

Higher order kinematics of [CII] emission line



対局的な回転成分を差し引くと速度残差に $m=2$ のパターンが残った。


Tsukui + 2023c in press

軸対称なinflow outflowでは対称な位置に反対符号の残差が現れるはず。


バー、渦巻き構造による非回転運動は $m=1$ or 3 のパターンが現れる。(Schoenmakers+97).

円盤の垂直方向の振動 (disk-bending wave, U-shaped warpping)を見ている。

Disk bending mode caused by interaction/gas accretion



Bland-Hawthorn and Topper-Garcia+21




Tsukui + 2023c in press

衛星銀河やガス降着による摂動によって円盤内部に $m=2$ の円盤振動を起動する。振動波は渦巻き腕と位置が一致する。


Galaxy transformation at $z=4.4$

Massive compact quiescent at $z > 2$


Newman+17




Bland-Hawthorn and Topper-Garcia+21





Dekel+09




Ito+23



(A) Velocity data



(D) Data - model odd term





Cold gas/satellite accretion and mergers supply gas in the **gas-rich gravitationally unstable** disk, causing the **seismic ripple of the disk (vertical disk bending wave)** and the **two-armed spiral**. Gas rich disk formed **star-forming gaseous bar**, subsequently triggering the **SF bulge (compact rotating core)** and **optical quasar**.

Evolutionally link between two populations

Massive compact quiescent?

Newman+17



Rizzo+2020


SFR $\sim 100 M_{\odot} \text{yr}^{-1}$

Gas depletion time = $\sim 360 \text{Myr}$


SFR $\sim 352 M_{\odot} \text{yr}^{-1}$

Gas depletion time $\sim 40 \text{Myr}$

Compaction?




Ito+23



Lelli+2021

SFR $\sim 1000 M_{\odot} \text{yr}^{-1}$

Gas depletion time $\sim 20 \text{Myr}$



Tsukui and Iguchi 2021


SFR $\sim 5000 M_{\odot} \text{yr}^{-1} (?)$

Gas depletion time $\sim 20 \text{Myr}$

Secular evolution?

Extended quenched spiral?

Fudamoto +2022




どのような条件でスターバーストが起き、遷移の過程で何が起きているか?

ALMA: ダストを見通し、空間分解した 1) 銀河構造、ダイナミクス and 2) 正確な星形成分布の把握

Star formation rate traced by FIR emission

- ダストが主に若い星で加熱されている。
- ダストに隠された星形成率は赤外線高度から見積もられる。 $SFR \propto L_{IR} \propto T_{dust}^{\sim 6}$




Herschel / Spitzer etc.

Young massive stars



Star formation rate traced by FIR emission

- ダストが主に若い星で加熱されている。
- ダストに隠された星形成率は赤外線高度から見積もられる。 $SFR \propto L_{IR} \propto T_{dust}^{\sim 6}$



Herschel / Spitzer etc.

UV light




Young massive stars




AGN

How much can AGN contribute to IR luminosity?

- 赤外光度が大きくなるほどAGNの赤外線光度に対する寄与は大きくなる(e.g., Alonso-Herrero+12 BPT/MIR spec diagnostics) up to ~100% for the bright source ($L_{\text{IR}} > 10^{13} L_{\odot}$)?(Symeonidis+21)
- AGNと星形成の分離は銀河全体の(空間分解していない)SEDによって行われてきた。
-> 仮定したテンプレートに依存する。(MIR: AGN torus model etc.).
- 遠赤外線波長 ($> \sim 100 \mu\text{m}$), AGNの寄与は無視できるとされてきた。が、



Alonso-Herrero et al. 2012




Leipski et al. 2014

AGN contribution in spatially resolved FIR image?


- 近年の高分解能ALMA dataは遠赤外線画像へのAGN寄与を示す。

J1243 z=7 3.6kpc resolution (Izumi+21)



J2348 z=7 200pc resolution (Walter+22)

$T_{\text{dust}} \sim > 200\text{K} ?$
SFR = $970 M_{\odot} \text{yr}^{-1}$ within $r < 110$ pc?
Continuum flux density ($\mu\text{Jy beam}^{-1}$)






- 1バンド観測のためダストの性質(温度、光学的厚み)に制限をつけることができない。

Star formation rate traced by FIR emission

- 空間分解してダスト温度を測定し、AGNの可能性を検証し、より正確な星形成率の把握を目指す。



ALMA observation





Young massive stars




AGN

Energetics of the disk interstellar medium


- Rest $161\mu\text{m}$ and $90\mu\text{m}$ はcoldダストのピークSEDに近い ($\sim 40\text{K}$),
フラックス比が温度の制限をつけられる (see also Shao et al. 2022).



Energetics of the disk interstellar medium





Tsukui + 2023b, MNRAS



温度ピーク $\sim 57 \text{ K} > 47\text{K}$ (typical quasar) $> 40\text{K}$ (typical star-forming galaxy)

温度ピークの位置は可視のクエーサーの位置 + FIR画像に存在する点源に一致する。

Energetics of the disk interstellar medium






温度ピーク $\sim 57 \text{ K} > 47\text{K}$ (typical quasar) $> 40\text{K}$ (typical star-forming galaxy)

温度ピークの位置は可視のクエーサーの位置 + FIR画像に存在する点源に一致する。

AGN-host galaxy decomposition

Whole galaxy's SED



Tsukui + 2023b, MNRAS



AGN-host galaxy decomposition




Whole galaxy's SED

Tsukui + 2023b, MNRAS





Point source

Host galaxy



Point source

Host galaxy




$$\text{updated SFR} = 1700^{+500}_{-400} M_{\odot} \text{yr}^{-1}$$

< 3x previous estimate $5000 M_{\odot} \text{yr}^{-1}$



due to a high AGN fraction $53^{+14}_{-15} \%$

AGN subtracted temperature in image space



Point source contributions to the total flux in the FIR bands are small $\sim 20\%$, but large $\sim 50\%$ in central pixel.


AGN subtracted temperature in image space



AGN component entirely responsible for the high dust temperature.

We underestimate the size of the SFR distribution if we neglect AGN

Kennicutt–Schmidt diagram



After AGN correction, BRI 1335-0417 is not an outlier but rather a normal starburst population with a gas depletion time of 100 Myr.

Summary

Spatially resolved ALMA data identify bulge, disk, spiral, and bar structure in BRI 1335-0417 at $z=4.4$

Higher order velocity residual in the disk indicates the vertical oscillation of the disk (U-shaped warping).

Gas or satellite accretion likely causes the seismic ripple and spiral arms in the disk.

Bar structure may channel the gas into the central quasar and bulge.

Assuming a single dust temperature over the galaxies leads to a false interpretation of the SFR, its size etc.

Without spatially resolved data, dust optical depth has no physical meaning: we link the opacity with dust mass given the physical beam area.


Open question

AGN-host decomposition will never be perfect:


AGN optical-UV emission may leak out from the dust torus and heat the extended region of galaxies.

(need more data, sample).

Galaxy transformation at $z=4.4$




Bland-Hawthorn and Topper-Garcia+21




Dekel+09

Massive compact quiescent at $z > 2$



Newman+17




Ito+23



(A) Velocity data




(D) Data - model odd term



Cold gas/satellite accretion and mergers supply gas in the **gas-rich gravitationally unstable disk**, causing the **seismic ripple of the disk (vertical disk bending wave)** and the **two-armed spiral**. Gas rich disk formed **star-forming gaseous bar**, subsequently triggering the **SF bulge (compact rotating core)** and **optical quasar**.

Fit optically-thick graybody function to each pixel (inc. CMB heating/contrast)

Free Parameters: T_{dust} and M_{dust} (or $\tau_{161\mu\text{m}}$) with $\beta = 2.14 \pm 0.17$




- First 2d temperature map without assuming an optically thin limit: linking dust optical depth with dust masses and physical area subtended by the beam.

AGN-host galaxy decomposition


Stardust code (Kokorev 2021), modified to accept point source as AGN constraint

With spatially resolved constraints



SFR~1700 Msun/yr

Without spatially resolved constraints




SFR~5260 Msun/yr

The goodness of fit does not change -> Only spatially integrated SED has no power to constrain the AGN fraction.

Identification of the spiral structure


We expand intensity image into logarithmic spiral models with a range of pitch angles (α) and number of arms (m).

Fourier expansion (basis function: sin cos)



Log spiral expansion (basis function: log spirals)
Puerari and Dottori 1992.



De-projected
Intensity image







$m=2, \alpha=33\text{deg}$ $m=2, \alpha=18\text{deg}$ $m=1, \alpha=33\text{deg}$ $m=2, \alpha=33\text{deg}$

We find that the dominant component is two-arm spirals with a pitch angle of 26.7deg. (still highest- z spiral structure)

$m=1$ corresponds to the lopsided structure.



Identification of bar structure



Tsukui + 2023c in press


Defining bar length maximum of the ellipticity profile (Wozniak and Pierce 1991)

Bar length $R_{\text{bar}} \sim 3.3$ kpc (cf. disk scale $R_{\text{disk}} \sim 1.83$ kpc) x2 bigger than nearby stellar bar length/disk size ratio.

The presence of a bar in gas-rich disk??


Stellar bar grows faster in a disk-dominated system (Fujii+18).

Gaseous disk may suppress stellar bar formation (Łokas 20; Athanassoula+13).



The experiment is done for isolated / a moderate gas mass fraction of up to 30%.


Gaseous bars may be prevalent in the gas-rich high-z universe.




Triaxial bar seen in dust continuum (Hodge+18; Gullberg+18).

The presence of a bar in gas-rich disk??

In gas-rich galaxies being actively supplied with gas, we may need a different perspective:
Gaseous bars form out of gravitationally unstable gas disks.




100% gaseous axisymmetric disk form stable gaseous bar



Cazes and Tohline+1999

The presence of a bar in gas-rich disk??

In gas-rich galaxies being actively supplied with gas, we may need a different perspective:
Gaseous bars form out of gravitationally unstable gas disks.





100% gaseous axisymmetric disk form stable gaseous bar

Controlled simulations focused on the gaseous bar with realistic star formation and gas accretion to address:

How long a gaseous bar can live?


The gaseous bar can yield a stellar bar later?

How much gaseous bar can help the starburst and AGN activity?




Cazes and Tohline+1999

Starforming bulge




Band7 continuum distribution




Bulge mass radius <1.3kpc is consistent with the size of the SF core ~800 pc

Image decomposition

- Band 7 image



- Band 9 image



A1. Dust temperature correlate with gas velocity dispersion?

- High dust temperature peaks preferentially aligned disk minor axis?
- [CII] velocity dispersion shows the conical enhanced structure.

